



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

MECHANICS.

153. Proposed by W. J. GREENSTREET, A. M., Editor of The Mathematical Gazette, Stroud, Eng.

An equiangular polygon consisting of equal, freely jointed rods, is hung up from vertex, A . The vertices adjacent to A are connected by a light rod of such length that the polygon is still regular. Find the stress in the rod and the reactions at the vertices.

Solution by G. B. M. ZERR, A. M., Ph. D., Professor of Chemistry and Physics, The Temple College, Philadelphia, Pa.

Let P be the point of suspension, O , the centroid of the system, T , the stress on the weightless rod FB , nW , the weight of the rods, $FB=x$, $AB=a$, $AO=y$, $\angle OAB=\angle OBA=\theta$, R =reaction.

Suppose the deformation such that the centroid moves vertically through a small space. Then $Tdx - nWdy = 0$.

$$BG : AB = \sin \theta : 1; \therefore x = 2a \sin \theta.$$

$$AO : AB = \sin \theta : \sin(\pi - 2\theta); \therefore y = a \sin \theta / \sin(\pi - 2\theta).$$

$$\therefore y = a \sin \theta / \sin 2\theta = \frac{1}{2} a \sec \theta.$$

$$\therefore dx = 2a \cos \theta d\theta, dy = \frac{1}{2} a \sec \theta \tan \theta d\theta.$$


$$\therefore 2a T \cos \theta d\theta = \frac{1}{2} (nW a) \sec \theta \tan \theta d\theta.$$

$$\therefore T = \frac{1}{4} n W \sec^2 \theta \tan \theta. \quad R \text{ acts along } BO, \angle ABG = \frac{1}{2} \pi - \theta.$$

$$\therefore \angle OBG = \angle OFG = 2\theta - \frac{1}{2} \pi, \angle BOF = 2\pi - 4\theta.$$

$$\therefore R : T = \sin(2\theta - \frac{1}{2} \pi) : \sin(2\pi - 4\theta); R : T = \cos 2\theta : \sin 4\theta.$$

$$\therefore R = T \cos 2\theta / \sin 4\theta = T / 2 \sin 2\theta = T / 4 \sin \theta \cos \theta. \quad \therefore R = \frac{1}{16} n W \sec^4 \theta.$$



Also solved by G. W. GREENWOOD.

154. Proposed by M. E. GRABER, Graduate Student, Heidelberg University, Tiffin, Ohio.

Find the form of the curve in a vertical plane, such that a heavy bar resting on its concave side and on a peg at a given point, (the origin), may be at rest at all positions.

Solution by G. W. GREENWOOD, A. B., Professor of Mathematics, McKendree College, Lebanon, Ill.; and E. L. SHERWOOD, Professor of Mathematics, Shady Side Academy, Pittsburgh, Pa.

Let the bar be homogeneous and of length $2b$. Assume the peg and the curve to be smooth. Take the peg as origin, the horizontal line through it in the plane as initial line, and measure θ downwards.

The coördinates of the lower extremity being (r, θ) , those of the center of the rod are $(r-b, \theta)$. Since in all positions the rod is in equilibrium, the center is at a constant distance from the initial line. Hence $(r-b) \sin \theta = a$, is the curve traced by one extremity.

Also solved by G. B. M. ZERR.

DIOPHANTINE ANALYSIS.

107. Proposed by L. C. WALKER, A. M., Graduate Student, Leland Stanford Jr. University, Cal.

Required the least three positive integral numbers such that the sum of all three of them and the sum of every two of them shall be a square number.